



Phase 3 Trial of Libtayo® (cemiplimab-rwlc) Combined with Chemotherapy Stopped Early Due to Significant Improvement in Overall Survival in Patients with First-line Advanced Non-small Cell Lung Cancer

August 5, 2021

UPDATE: This press release has been updated to state that progression-free survival (PFS) is a secondary endpoint in the trial. In the "About the Phase 3 Trial" section of the previous version, PFS was listed as a co-primary endpoint.

TARRYTOWN, N.Y. and PARIS, Aug. 5, 2021 /PRNewswire/ --

Libtayo combined with chemotherapy increased median overall survival from 13 to 22 months, leading to a 29% reduction in the risk of death

Trial enrolled patients with locally advanced and metastatic disease with squamous or non-squamous histology, across all PD-L1 expression levels

Libtayo has now demonstrated improved overall survival as a monotherapy or in combination with chemotherapy in first-line advanced non-small cell lung cancer

Regeneron Pharmaceuticals, Inc. (NASDAQ: **REGN**) and Sanofi today announced the Phase 3 trial of PD-1 inhibitor Libtayo in combination with platinum-doublet chemotherapy was stopped early after meeting its overall survival (OS) primary endpoint in patients with advanced non-small cell lung cancer (NSCLC). Adding Libtayo to chemotherapy significantly improved OS, compared to chemotherapy alone, in the trial that enrolled patients with metastatic or locally advanced disease and tumors with either squamous or non-squamous histology and across all PD-L1 expression levels. These data are planned to form the basis of regulatory submissions in the U.S. and European Union.

"Libtayo in combination with chemotherapy increased median overall survival to 22 months in patients with advanced non-small cell lung cancer, compared to 13 months with chemotherapy alone," said Miranda Gogishvili, M.D., an oncologist at the High Technology Medical Center, University Clinic, in Tbilisi, Georgia and a trial investigator. "Notably, the Phase 3 trial enrolled patients with a variety of challenging-to-treat disease characteristics, as well as those with locally advanced disease. These data add to the growing body of evidence supporting Libtayo in advanced non-small cell lung cancer, which also include the pivotal results for Libtayo monotherapy in cases of high PD-L1 expression."

The decision to stop the trial early was based on a recommendation by the Independent Data Monitoring Committee (IDMC) during a protocol-specified interim analysis. In this top-line initial analysis of 466 patients, combining Libtayo with chemotherapy reduced the risk of death by 29% compared to chemotherapy alone (hazard ratio: 0.71; 95% confidence interval [CI]: 0.53-0.93; p=0.014). Median OS was 22 months (95% CI: 16 months to not evaluable) for Libtayo and chemotherapy, and 13 months (95% CI: 12 to 16 months) for chemotherapy alone. No new Libtayo safety signals were identified in the IDMC analysis, and additional detailed efficacy and safety data will be presented at an upcoming medical meeting.

Lung cancer is the leading cause of cancer death worldwide. In 2020, an estimated 2.2 million and 225,000 new cases were diagnosed globally and in the U.S., respectively. Approximately 84% of all lung cancers are NSCLC, with 75% of these cases diagnosed in advanced stages. While PD-1 inhibitor monotherapy has primarily advanced the treatment of NSCLC with ≥50% PD-L1 expression, approximately 70% of all NSCLC cases will have <50% PD-L1 expression, making it the most common treatment setting.

The use of Libtayo in combination with chemotherapy for advanced NSCLC is currently under clinical investigation, and its safety and efficacy have not been fully evaluated by any regulatory authority.

About the Phase 3 Trial

The randomized, multicenter Phase 3 trial, called EMPOWER-Lung 3, investigated a first-line combination treatment of Libtayo and platinum-doublet chemotherapy, compared to platinum-doublet chemotherapy alone, in squamous or non-squamous advanced NSCLC irrespective of PD-L1 expression. Specifically, the trial included 466 patients who tested negative for ALK, EGFR and ROS1 mutations and had either previously untreated metastatic NSCLC (stage IV) or locally advanced NSCLC (stage IIIB/C) and were not candidates for definitive chemoradiation.

Patients were randomized 2:1 to receive either Libtayo 350 mg (n=312) or placebo (n=154) administered intravenously every three weeks for 108 weeks, plus platinum-doublet chemotherapy administered every three weeks for four cycles. The primary endpoint was OS, and secondary endpoints included progression-free survival, objective response rate and best overall response.

Among trial patients, 30% (n=139) had tumors with <1% PD-L1 expression, 38% (n=175) had tumors with 1% to 49% PD-L1 expression, and 33% (n=152) had tumors with ≥50% PD-L1 expression.

About Libtayo

Libtayo is a fully human monoclonal antibody targeting the immune checkpoint receptor PD-1 on T-cells. By binding to PD-1, Libtayo has been shown to block cancer cells from using the PD-1 pathway to suppress T-cell activation.

The generic name for Libtayo in its approved U.S. indications is cemiplimab-rwlc, with rwlc as the suffix designated in accordance with Nonproprietary Naming of Biological Products Guidance for Industry issued by the U.S. FDA. Libtayo is being jointly developed by Regeneron and Sanofi under a global collaboration agreement.

The extensive clinical program for Libtayo is focused on difficult-to-treat cancers. Libtayo is currently being investigated in advanced cervical cancer,

as well as in trials combining Libtayo with either conventional or novel therapeutic approaches for other solid tumors and blood cancers. These potential uses are investigational, and their safety and efficacy have not been evaluated by any regulatory authority.

About Regeneron's *VelocImmune*[®] Technology

Regeneron's *VelocImmune* technology utilizes a proprietary genetically engineered mouse platform endowed with a genetically humanized immune system to produce optimized fully human antibodies. When Regeneron's President and Chief Scientific Officer George D. Yancopoulos was a graduate student with his mentor Frederick W. Alt in 1985, they were the first to [envision](#) making such a genetically humanized mouse, and Regeneron has spent decades inventing and developing *VelocImmune* and related *VelociSuite*[®] technologies. Dr. Yancopoulos and his team have used *VelocImmune* technology to create approximately a quarter of all original, FDA-approved fully human monoclonal antibodies currently available. This includes REGEN-COV[™] (casirivimab and imdevimab), Dupixent[®] (dupilumab), Libtayo (cemiplimab-rwlc), Praluent[®] (alirocumab), Kevzara[®] (sarilumab), Evkeeza[®] (evinacumab-dgnb) and Imzab[™] (atoltivimab, maftivimab and odesivimab-ebgn).

IMPORTANT SAFETY INFORMATION AND INDICATION FOR U.S. PATIENTS

What is Libtayo?

Libtayo is a prescription medicine used to treat people with a type of skin cancer called cutaneous squamous cell carcinoma (CSCC) that has spread or cannot be cured by surgery or radiation.

Libtayo is a prescription medicine used to treat people with a type of skin cancer called basal cell carcinoma that cannot be removed by surgery (locally advanced BCC) and have received treatment with an HHI, or cannot receive treatment with a HHI.

Libtayo is a prescription medicine used to treat people with a type of skin cancer called basal cell carcinoma that has spread (metastatic BCC) and have received treatment with a hedgehog pathway inhibitor (HHI), or cannot receive treatment with a HHI. This use is approved based on how many patients responded to treatment and how long they responded. Studies are ongoing to provide additional information about clinical benefit.

Libtayo is a prescription medicine used to treat people with a type of lung cancer called non-small cell lung cancer (NSCLC). Libtayo may be used as your first treatment when your lung cancer has not spread outside your chest (locally advanced lung cancer) and you cannot have surgery or chemotherapy with radiation, or your lung cancer has spread to other areas of your body (metastatic lung cancer), and your tumor tests positive for high "PD-L1" and your tumor does not have an abnormal "EGFR," "ALK" or "ROS1" gene.

It is not known if Libtayo is safe and effective in children.

What is the most important information I should know about Libtayo?

Libtayo is a medicine that may treat certain cancers by working with your immune system. Libtayo can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms, including:

- **Lung problems:** cough, shortness of breath, or chest pain
- **Intestinal problems:** diarrhea (loose stools) or more frequent bowel movements than usual, stools that are black, tarry, sticky or have blood or mucus, or severe stomach-area (abdomen) pain or tenderness
- **Liver problems:** yellowing of your skin or the whites of your eyes, severe nausea or vomiting, pain on the right side of your stomach area (abdomen), dark urine (tea colored), or bleeding or bruising more easily than normal
- **Hormone gland problems:** headache that will not go away or unusual headaches, eye sensitivity to light, eye problems, rapid heartbeat, increased sweating, extreme tiredness, weight gain or weight loss, feeling more hungry or thirsty than usual, urinating more often than usual, hair loss, feeling cold, constipation, your voice gets deeper, dizziness or fainting, or changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness
- **Kidney problems:** decrease in your amount of urine, blood in your urine, swelling of your ankles, or loss of appetite
- **Skin problems:** rash, itching, skin blistering or peeling, painful sores or ulcers in mouth or nose, throat, or genital area, fever or flu-like symptoms, or swollen lymph nodes
- **Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with Libtayo. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:** chest pain, irregular heartbeat, shortness of breath or swelling of ankles, confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs, double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight, persistent or severe muscle pain or weakness, muscle cramps, low red blood cells, or bruising
- **Infusion reactions that can sometimes be severe.** Signs and symptoms of infusion reactions may include: nausea, chills or shaking, itching or rash, flushing, shortness of breath or wheezing, dizziness, feel like passing out, fever, back or neck pain, or facial swelling.
- **Rejection of a transplanted organ.** Your healthcare provider should tell you what signs and symptoms you should report and monitor you, depending on the type of organ transplant that you have had.
- **Complications, including graft-versus-host disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).** These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with Libtayo. Your healthcare provider will monitor you for these complications.

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with Libtayo. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may also need to delay or completely stop treatment with Libtayo if you have severe side effects.

Before you receive Libtayo, tell your healthcare provider about all your medical conditions, including if you:

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. Libtayo can harm your unborn baby

Females who are able to become pregnant:

- - Your healthcare provider will give you a pregnancy test before you start treatment.
 - You should use an effective method of birth control during your treatment and for at least 4 months after your last dose of Libtayo. Talk with your healthcare provider about birth control methods that you can use during this time.
 - Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with Libtayo.
- are breastfeeding or plan to breastfeed. It is not known if Libtayo passes into your breast milk.
Do not breastfeed during treatment and for at least 4 months after the last dose of Libtayo.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of Libtayo include muscle or bone pain, tiredness, rash, and diarrhea. These are not all the possible side effects of Libtayo. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Regeneron Pharmaceuticals and Sanofi at 1-877-542-8296.

Please see accompanying full [Prescribing Information](#), including [Medication Guide](#).

About Regeneron

Regeneron (NASDAQ: REGN) is a leading biotechnology company that invents life-transforming medicines for people with serious diseases. Founded and led for over 30 years by physician-scientists, our unique ability to repeatedly and consistently translate science into medicine has led to nine FDA-approved treatments and numerous product candidates in development, almost all of which were homegrown in our laboratories. Our medicines and pipeline are designed to help patients with eye diseases, allergic and inflammatory diseases, cancer, cardiovascular and metabolic diseases, pain, hematologic conditions, infectious diseases and rare diseases.

Regeneron is accelerating and improving the traditional drug development process through our proprietary *VelociSuite* technologies, such as *VelocImmune*, which uses unique genetically humanized mice to produce optimized fully human antibodies and bispecific antibodies, and through ambitious research initiatives such as the Regeneron Genetics Center, which is conducting one of the largest genetics sequencing efforts in the world.

For additional information about the company, please visit www.regeneron.com or follow @Regeneron on Twitter.

About Sanofi

Sanofi is dedicated to supporting people through their health challenges. We are a global biopharmaceutical company focused on human health. We prevent illness with vaccines, provide innovative treatments to fight pain and ease suffering. We stand by the few who suffer from rare diseases and the millions with long-term chronic conditions.

With more than 100,000 people in 100 countries, Sanofi is transforming scientific innovation into healthcare solutions around the globe.

Regeneron Forward-Looking Statements and Use of Digital Media

This press release includes forward-looking statements that involve risks and uncertainties relating to future events and the future performance of Regeneron Pharmaceuticals, Inc. ("Regeneron" or the "Company"), and actual events or results may differ materially from these forward-looking statements. Words such as "anticipate," "expect," "intend," "plan," "believe," "seek," "estimate," variations of such words, and similar expressions are intended to identify such forward-looking statements, although not all forward-looking statements contain these identifying words. These statements concern, and these risks and uncertainties include, among others, the impact of SARS-CoV-2 (the virus that has caused the COVID-19 pandemic) on Regeneron's business and its employees, collaborators, and suppliers and other third parties on which Regeneron relies, Regeneron's and its collaborators' ability to continue to conduct research and clinical programs, Regeneron's ability to manage its supply chain, net product sales of products marketed or otherwise commercialized by Regeneron and/or its collaborators (collectively, "Regeneron's Products"), and the global economy; the nature, timing, and possible success and therapeutic applications of Regeneron's Products and product candidates being developed by Regeneron and/or its collaborators (collectively, "Regeneron's Product Candidates") and research and clinical programs now underway or planned, including without limitation Libtayo® (cemiplimab) in combination with chemotherapy for the treatment of patients with advanced non-small cell lung cancer ("NSCLC"); the likelihood, timing, and scope of possible regulatory approval and commercial launch of Regeneron's Product Candidates and new indications for Regeneron's Products, such as possible regulatory approval of Libtayo in combination with chemotherapy for NSCLC and as monotherapy for advanced cervical cancer (as well as in combination with either conventional or novel therapeutic approaches for both solid tumors and blood cancers); uncertainty of the utilization, market acceptance, and commercial success of Regeneron's Products (such as Libtayo) and Regeneron's Product Candidates and the impact of studies (whether conducted by Regeneron or others and whether mandated or voluntary), including the study discussed in this press release, on any of the foregoing; the ability of Regeneron's collaborators, suppliers, or other third parties (as applicable) to perform manufacturing, filling, finishing, packaging, labeling, distribution, and other steps related to Regeneron's Products and Regeneron's Product Candidates; the ability of Regeneron to manufacture and manage supply chains for multiple products and product candidates; safety issues resulting from the administration of Regeneron's Products (such as Libtayo) and Regeneron's Product Candidates in patients, including

serious complications or side effects in connection with the use of Regeneron's Products and Regeneron's Product Candidates in clinical trials; determinations by regulatory and administrative governmental authorities which may delay or restrict Regeneron's ability to continue to develop or commercialize Regeneron's Products and Regeneron's Product Candidates, including without limitation Litbayo; ongoing regulatory obligations and oversight impacting Regeneron's Products, research and clinical programs, and business, including those relating to patient privacy; the availability and extent of reimbursement of Regeneron's Products from third-party payers, including private payer healthcare and insurance programs, health maintenance organizations, pharmacy benefit management companies, and government programs such as Medicare and Medicaid; coverage and reimbursement determinations by such payers and new policies and procedures adopted by such payers; competing drugs and product candidates that may be superior to, or more cost effective than, Regeneron's Products and Regeneron's Product Candidates; the extent to which the results from the research and development programs conducted by Regeneron and/or its collaborators may be replicated in other studies and/or lead to advancement of product candidates to clinical trials, therapeutic applications, or regulatory approval; unanticipated expenses; the costs of developing, producing, and selling products; the ability of Regeneron to meet any of its financial projections or guidance and changes to the assumptions underlying those projections or guidance; the potential for any license, collaboration, or supply agreement, including Regeneron's agreements with Sanofi, Bayer, and Teva Pharmaceutical Industries Ltd. (or their respective affiliated companies, as applicable), to be cancelled or terminated; and risks associated with intellectual property of other parties and pending or future litigation relating thereto (including without limitation the patent litigation and other related proceedings relating to EYLEA[®] (afibercept) Injection, Dupixent[®] (dupilumab), Praluent[®] (alirocumab), and REGEN-COV[™] (casirivimab and imdevimab)), other litigation and other proceedings and government investigations relating to the Company and/or its operations, the ultimate outcome of any such proceedings and investigations, and the impact any of the foregoing may have on Regeneron's business, prospects, operating results, and financial condition. A more complete description of these and other material risks can be found in Regeneron's filings with the U.S. Securities and Exchange Commission. Any forward-looking statements are made based on management's current beliefs and judgment, and the reader is cautioned not to rely on any forward-looking statements made by Regeneron. Regeneron does not undertake any obligation to update (publicly or otherwise) any forward-looking statement, including without limitation any financial projection or guidance, whether as a result of new information, future events, or otherwise.

Regeneron uses its media and investor relations website and social media outlets to publish important information about the Company, including information that may be deemed material to investors. Financial and other information about Regeneron is routinely posted and is accessible on Regeneron's media and investor relations website (<http://newsroom.regeneron.com>) and its Twitter feed (<http://twitter.com/regeneron>).

Sanofi Forward-Looking Statements

This press release contains forward-looking statements as defined in the Private Securities Litigation Reform Act of 1995, as amended. Forward-looking statements are statements that are not historical facts. These statements include projections and estimates and their underlying assumptions, statements regarding plans, objectives, intentions and expectations with respect to future financial results, events, operations, services, product development and potential, and statements regarding future performance. Forward-looking statements are generally identified by the words "expects", "anticipates", "believes", "intends", "estimates", "plans" and similar expressions. Although Sanofi's management believes that the expectations reflected in such forward-looking statements are reasonable, investors are cautioned that forward-looking information and statements are subject to various risks and uncertainties, many of which are difficult to predict and generally beyond the control of Sanofi, that could cause actual results and developments to differ materially from those expressed in, or implied or projected by, the forward-looking information and statements. These risks and uncertainties include among other things, the uncertainties inherent in research and development, future clinical data and analysis, including post marketing, decisions by regulatory authorities, such as the FDA or the EMA, regarding whether and when to approve any drug, device or biological application that may be filed for any such product candidates as well as their decisions regarding labelling and other matters that could affect the availability or commercial potential of such product candidates, the fact that product candidates if approved may not be commercially successful, the future approval and commercial success of therapeutic alternatives, Sanofi's ability to benefit from external growth opportunities, to complete related transactions and/or obtain regulatory clearances, risks associated with intellectual property and any related pending or future litigation and the ultimate outcome of such litigation, trends in exchange rates and prevailing interest rates, volatile economic and market conditions, cost containment initiatives and subsequent changes thereto, and the impact that COVID-19 will have on us, our customers, suppliers, vendors, and other business partners, and the financial condition of any one of them, as well as on our employees and on the global economy as a whole. Any material effect of COVID-19 on any of the foregoing could also adversely impact us. This situation is changing rapidly and additional impacts may arise of which we are not currently aware and may exacerbate other previously identified risks. The risks and uncertainties also include the uncertainties discussed or identified in the public filings with the SEC and the AMF made by Sanofi, including those listed under "Risk Factors" and "Cautionary Statement Regarding Forward-Looking Statements" in Sanofi's annual report on Form 20-F for the year ended December 31, 2020. Other than as required by applicable law, Sanofi does not undertake any obligation to update or revise any forward-looking information or statements.

Regeneron Contacts:

Media Relations

Daren Kwok
Tel: +1 914-847-1328
daren.kwok@regeneron.com

Investor Relations

Vesna Tosic
Tel: +1 914-847-5443
vesna.tosic@regeneron.com

Sanofi Contacts:

Media Relations

Sally Bain
Tel: +1 781-264-1091
sally.bain@sanofi.com

Investor Relations Paris

Eva Schaefer-Jansen
Arnaud Delepine
Nathalie Pham

Investor Relations North America

Felix Lauscher
Fara Berkowitz
Suzanne Greco

IR main line:
Tel: +33 (0)1 53 77 45 45
investor.relations@sanofi.com
<https://www.sanofi.com/en/investors/contact>

 View original content: <https://www.prnewswire.com/news-releases/phase-3-trial-of-libtayo-cemiplimab-nwlc-combined-with-chemotherapy-stopped-early-due-to-significant-improvement-in-overall-survival-in-patients-with-first-line-advanced-non-small-cell-lung-cancer-301348999.html>

SOURCE Regeneron Pharmaceuticals, Inc.