



Dupixent® (dupilumab) Late-breaking Phase 3 COPD Results Presented at ATS and Simultaneously Published in the New England Journal of Medicine

May 21, 2023

Dupixent is the first and only investigational biologic for COPD that has demonstrated a significant reduction in moderate or severe acute exacerbations by 30% compared to placebo

Dupixent is the first and only investigational biologic for COPD that has significantly improved lung function at 12 and 52 weeks, with numerical improvements seen as early as 2 weeks

Dupixent significantly improved quality of life, with numerical improvements as early as 4 weeks after initiating treatment, and respiratory symptoms

COPD is the third leading cause of death worldwide, with no new treatment approaches approved in more than a decade; trial enrolled patients with moderate-to-severe disease and evidence of type 2 inflammation (i.e., blood eosinophils ≥ 300 cells/ μ L)

TARRYTOWN, N.Y. and PARIS, May 21, 2023 (GLOBE NEWSWIRE) -- Regeneron Pharmaceuticals, Inc. (NASDAQ: REGN) and Sanofi today presented positive Phase 3 results evaluating the investigational use of Dupixent® (dupilumab) compared to placebo in adults currently on maximal standard-of-care inhaled therapy (triple therapy) with uncontrolled chronic obstructive pulmonary disease (COPD) and evidence of type 2 inflammation. The Dupixent Phase 3 results were presented in the 2023 American Thoracic Society (ATS) International Conference session “New England Journal of Medicine and JAMA. Discussion on the Edge: Reports of Recently Published Pulmonary Research” and simultaneously published in the [New England Journal of Medicine \(NEJM\)](#). These results will also be presented in the “Breaking News: Clinical Trial Results in Pulmonary Medicine” session on May 22.

“I’ve seen patients with uncontrolled chronic obstructive pulmonary disease struggle for far too long with the debilitating symptoms of this progressive disease – with limited, incremental improvement on current treatment options,” said Surya Bhatt, M.D., MSPH, Associate Professor at the University of Alabama at Birmingham Division of Pulmonary, Allergy, and Critical Care Medicine, and a co-principal investigator of the trial. “This trial showed that dupilumab has the potential to impact the vicious cycle of exacerbations and lung function decline in patients with uncontrolled COPD with type 2 inflammation, and significantly improve respiratory symptoms. Dupilumab also helped improve health-related quality of life measures, which, from my years of experience as a physician, are just as meaningful for patients as being able to breathe easier.”

COPD is a life-threatening respiratory disease that damages the lungs and causes progressive lung function decline. Symptoms include persistent cough and breathlessness that may not only impair the ability to perform routine daily activities, but can also lead to anxiety, depression and sleep disturbances. COPD is also associated with a significant health and economic burden due to recurrent acute exacerbations that require systemic corticosteroid treatment and/or lead to hospitalization or even death. Smoking and exposure to noxious particles are key risk factors for COPD, but even individuals who quit smoking can still develop or continue having the disease. In the U.S. alone, approximately 300,000 people live with uncontrolled COPD with evidence of type 2 inflammation.

The results presented at ATS and published in *NEJM* are from the BOREAS trial, which [met](#) the primary and all key secondary endpoints. As presented and published, patients receiving Dupixent (n=468) compared to placebo (n=471) added to maximal standard-of-care inhaled triple therapy experienced a:

- 30% reduction in moderate or severe acute COPD exacerbations over 52 weeks ($p < 0.001$), the primary endpoint.
- 160 mL improvement in lung function from baseline at 12 weeks versus 77 mL ($p < 0.001$).
 - Numerical improvements were observed as early as 2 weeks, with the benefit versus placebo sustained through 52 weeks (Dupixent: 153 mL, placebo: 70 mL; $p < 0.001$).
- 9.7-point improvement in health-related quality of life (QoL; patient-reported outcome on a scale from 0-100) from baseline at 52 weeks versus a 6.4-point improvement ($p = 0.002$), with numerical improvements observed as early as 4 weeks.
- 2.7-point reduction in respiratory symptom severity (patient-reported outcome on a scale from 0-40) from baseline at 52 weeks versus a 1.6-point reduction ($p = 0.001$).

In a pre-specified analysis from a subgroup of patients (Dupixent n=195, placebo n=188) with elevated levels (≥ 20 ppb) of fractional exhaled nitric oxide (FeNO) – an airway biomarker of type 2 inflammation – Dupixent treatment also led to a significant 38% reduction in exacerbations compared to placebo at 52 weeks ($p = 0.005$). In this subgroup, Dupixent also led to an improvement in lung function of 232 mL versus 108 mL for placebo at 12 weeks ($p = 0.002$) that was sustained at 52 weeks with an improvement in lung function of 247 mL versus 120 mL for placebo ($p = 0.003$).

The safety results were generally consistent with the known safety profile of Dupixent in its approved indications. Overall rates of adverse events (AEs) were 77% for Dupixent and 76% for placebo. AEs more commonly observed with Dupixent compared to placebo included headache (8.1% Dupixent, 6.8% placebo), diarrhea (5.3% Dupixent, 3.6% placebo) and back pain (5.1% Dupixent, 3.4% placebo). AEs more commonly observed with placebo compared to Dupixent included upper respiratory tract infection (9.8% placebo, 7.9% Dupixent), hypertension (6.0% placebo, 3.6% Dupixent) and COVID-19 (5.7% placebo, 4.1% Dupixent). AEs leading to deaths were balanced between the two arms (1.7% placebo, 1.5% Dupixent).

The second, replicate Phase 3 trial of Dupixent in COPD with evidence of type 2 inflammation (NOTUS) is ongoing, with data expected in 2024. The safety and efficacy of Dupixent in COPD are currently under clinical investigation and have not been evaluated by any regulatory authority. Regeneron

and Sanofi look forward to discussing the BOREAS data with regulators.

About the Dupixent COPD Phase 3 Trial Program

BOREAS is one of two pivotal trials in the Dupixent COPD program. The randomized, Phase 3, double-blind, placebo-controlled trial evaluated the efficacy and safety of Dupixent in 939 adults who were current or former smokers aged 40 to 80 years with moderate-to-severe COPD. All patients in the trial had evidence of type 2 inflammation, as measured by blood eosinophils ≥ 300 cells/ μL . Patients with a diagnosis or history of asthma were excluded from the trial. During the 52-week treatment period, patients received Dupixent or placebo every two weeks added to a maximal standard-of-care inhaled triple therapy of inhaled corticosteroids (ICS), long-acting beta agonists (LABA), and long-acting muscarinic antagonists (LAMA). Double maintenance therapy, which included LABA and LAMA, was allowed if ICS was contraindicated.

The primary endpoint evaluated the annualized rate of acute moderate or severe COPD exacerbations. Moderate exacerbations were defined as those requiring systemic steroids and/or antibiotics. Severe exacerbations were defined as those: requiring hospitalization; requiring more than a day of observation in an emergency department or urgent care facility; or resulting in death.

Key secondary and other hierarchy endpoints included:

- Change from baseline in lung function (assessed by pre-bronchodilator forced expiratory volume over one second [FEV₁]) at 12 and 52 weeks in both the overall population and those with FeNO ≥ 20 ppb.
- Change from baseline at 52 weeks in St. George's Respiratory Questionnaire (SGRQ) total score compared to placebo (scale from 0-100).
- Change from baseline at 52 weeks in the Evaluating Respiratory Symptoms in COPD (E-RS: COPD) scale score (scale from 0-40).
- The annualized rate of acute moderate or severe COPD exacerbations in patients with FeNO ≥ 20 ppb.

About Regeneron and Sanofi's COPD Clinical Research Program

Regeneron and Sanofi are motivated to transform the treatment paradigm of COPD by examining the role different types of inflammation play in the disease progression through the investigation of two potentially first-in-class biologics, Dupixent and itepekimab.

Dupixent inhibits the signaling of the interleukin-4 (IL-4) and interleukin-13 (IL-13) pathways and the program focuses on a specific population of people with evidence of type 2 inflammation. Itepekimab is a fully human monoclonal antibody that binds to and inhibits interleukin-33 (IL-33), an initiator and amplifier of broad inflammation in COPD. Across both programs, four Phase 3 trials are ongoing and designed to inform next-generation treatments for people with COPD who might not have other options.

Itepekimab is currently under clinical investigation and its safety and efficacy have not been evaluated by any regulatory authority.

About Dupixent

Dupixent, which was invented using Regeneron's proprietary *VelocImmune*[®] technology, is a fully human monoclonal antibody that inhibits the signaling of the interleukin-4 (IL-4) and interleukin-13 (IL-13) pathways and is not an immunosuppressant. The Dupixent development program has shown significant clinical benefit and a decrease in type 2 inflammation in Phase 3 trials, establishing that IL-4 and IL-13 are key and central drivers of the type 2 inflammation that plays a major role in multiple related and often co-morbid diseases. These diseases include approved indications for Dupixent, such as atopic dermatitis, asthma, chronic rhinosinusitis with nasal polyposis (CRSwNP), eosinophilic esophagitis (EoE) and prurigo nodularis.

Dupixent has received regulatory approvals in one or more countries around the world for use in certain patients with atopic dermatitis, asthma, CRSwNP, EoE or prurigo nodularis in different age populations. Dupixent is currently approved for one or more of these indications in more than 60 countries, including in Europe, the U.S. and Japan. More than 600,000 patients are being treated with Dupixent globally.

About Regeneron's *VelocImmune* Technology

Regeneron's *VelocImmune* technology utilizes a proprietary genetically engineered mouse platform endowed with a genetically humanized immune system to produce optimized fully human antibodies. When Regeneron's co-Founder, President and Chief Scientific Officer George D. Yancopoulos was a graduate student with his mentor Frederick W. Alt in 1985, they were the first to [envision](#) making such a genetically humanized mouse, and Regeneron has spent decades inventing and developing *VelocImmune* and related *VelociSuite*[®] technologies. Dr. Yancopoulos and his team have used *VelocImmune* technology to create a substantial proportion of all original, FDA-approved or authorized fully human monoclonal antibodies. This includes REGEN-COV[®] (casirivimab and imdevimab), Dupixent, Libtayo[®] (cemiplimab-rwlc), Praluent[®] (alirocumab), Kevzara[®] (sarilumab), Evkeeza[®] (evinacumab-dgng) and Inmazeb[®] (atoltivimab, maftivimab and odesivimab-ebgn).

Dupilumab Development Program

Dupilumab is being jointly developed by Regeneron and Sanofi under a global collaboration agreement. To date, dupilumab has been studied across more than 60 clinical trials involving more than 10,000 patients with various chronic diseases driven in part by type 2 inflammation.

In addition to the currently approved indications, Regeneron and Sanofi are studying dupilumab in a broad range of diseases driven by type 2 inflammation or other allergic processes in Phase 3 trials, including pediatric EoE, chronic spontaneous urticaria, chronic pruritus of unknown origin, chronic obstructive pulmonary disease with evidence of type 2 inflammation and bullous pemphigoid. These potential uses of dupilumab are currently under clinical investigation, and the safety and efficacy in these conditions have not been fully evaluated by any regulatory authority.

U.S. INDICATIONS

DUPIXENT is a prescription medicine used:

- to treat adults and children 6 months of age and older with moderate-to-severe eczema (atopic dermatitis or AD) that is not well controlled with prescription therapies used on the skin (topical), or who cannot use topical therapies. DUPIXENT can be used with or without topical corticosteroids. It is not known if DUPIXENT is safe and effective in children with atopic

dermatitis under 6 months of age.

- with other asthma medicines for the maintenance treatment of moderate-to-severe eosinophilic or oral steroid dependent asthma in adults and children 6 years of age and older whose asthma is not controlled with their current asthma medicines. DUPIXENT helps prevent severe asthma attacks (exacerbations) and can improve your breathing. DUPIXENT may also help reduce the amount of oral corticosteroids you need while preventing severe asthma attacks and improving your breathing. DUPIXENT is not used to treat sudden breathing problems. It is not known if DUPIXENT is safe and effective in children with asthma under 6 years of age.
- with other medicines for the maintenance treatment of chronic rhinosinusitis with nasal polyposis (CRSwNP) in adults whose disease is not controlled. It is not known if DUPIXENT is safe and effective in children with chronic rhinosinusitis with nasal polyposis under 18 years of age.
- to treat adults and children 12 years of age and older, who weigh at least 88 pounds (40 kg), with eosinophilic esophagitis (EoE). It is not known if DUPIXENT is safe and effective in children with eosinophilic esophagitis under 12 years of age and who weigh at least 88 pounds (40 kg).
- to treat adults with prurigo nodularis (PN). It is not known if DUPIXENT is safe and effective in children with prurigo nodularis under 18 years of age.

IMPORTANT SAFETY INFORMATION

Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT®.

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- have eye problems.
- have a parasitic (helminth) infection.
- are scheduled to receive any vaccinations. You should not receive a “live vaccine” right before and during treatment with DUPIXENT.
- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.
 - A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby. To enroll or get more information call 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/dupixent/>.
- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines; have asthma and use an asthma medicine; or have atopic dermatitis, chronic rhinosinusitis with nasal polyposis, eosinophilic esophagitis, or prurigo nodularis and also have asthma.

Do not change or stop your corticosteroid medicine or other asthma medicine without talking to your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine or other asthma medicine to come back.

DUPIXENT can cause serious side effects, including:

- **Allergic reactions.** DUPIXENT can cause allergic reactions that can sometimes be severe. Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, joint pain, general ill feeling, itching, skin rash, swollen lymph nodes, nausea or vomiting, or cramps in your stomach-area.
- **Eye problems.** Tell your healthcare provider if you have any new or worsening eye problems, including eye pain or changes in vision, such as blurred vision. Your healthcare provider may send you to an ophthalmologist for an exam if needed.
- **Inflammation of your blood vessels.** Rarely, this can happen in people with asthma who receive DUPIXENT. This may happen in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered. It is not known whether this is caused by DUPIXENT. Tell your healthcare provider right away if you have: rash, chest pain, worsening shortness of breath, a feeling of pins and needles or numbness of your arms or legs, or persistent fever.
- **Joint aches and pain.** Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

The most common side effects include:

- **Eczema:** injection site reactions, eye and eyelid inflammation, including redness, swelling, and itching, sometimes with blurred vision, cold sores in your mouth or on your lips, and high count of a certain white blood cell (eosinophilia).

- **Asthma:** injection site reactions, high count of a certain white blood cell (eosinophilia), pain in the throat (oropharyngeal pain), and parasitic (helminth) infections.
- **Chronic Rhinosinusitis with Nasal Polyposis:** injection site reactions, eye and eyelid inflammation, including redness, swelling, and itching, sometimes with blurred vision, high count of a certain white blood cell (eosinophilia), gastritis, joint pain (arthralgia), trouble sleeping (insomnia), and toothache.
- **Eosinophilic Esophagitis:** injection site reactions, upper respiratory tract infections, cold sores in your mouth or on your lips, and joint pain (arthralgia).
- **Prurigo Nodularis:** eye and eyelid inflammation, including redness, swelling, and itching, sometimes with blurred vision, herpes virus infections, common cold symptoms (nasopharyngitis), dizziness, muscle pain, and diarrhea.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 6 months to less than 12 years of age, DUPIXENT should be given by a caregiver.

Please see accompanying full [Prescribing Information](#) including Patient Information.

About Regeneron

Regeneron (NASDAQ: REGN) is a leading biotechnology company that invents, develops and commercializes life-transforming medicines for people with serious diseases. Founded and led for 35 years by physician-scientists, our unique ability to repeatedly and consistently translate science into medicine has led to nine FDA-approved treatments and numerous product candidates in development, almost all of which were homegrown in our laboratories. Our medicines and pipeline are designed to help patients with eye diseases, allergic and inflammatory diseases, cancer, cardiovascular and metabolic diseases, pain, hematologic conditions, infectious diseases and rare diseases.

Regeneron is accelerating and improving the traditional drug development process through our proprietary *VelociSuite*[®] technologies, such as *VelocImmune*[®], which uses unique genetically humanized mice to produce optimized fully human antibodies and bispecific antibodies, and through ambitious research initiatives such as the Regeneron Genetics Center, which is conducting one of the largest genetics sequencing efforts in the world.

For more information, please visit www.Regeneron.com or follow @Regeneron on Twitter.

About Sanofi

We are an innovative global healthcare company, driven by one purpose: we chase the miracles of science to improve people's lives. Our team, across some 100 countries, is dedicated to transforming the practice of medicine by working to turn the impossible into the possible. We provide potentially life-changing treatment options and life-saving vaccine protection to millions of people globally, while putting sustainability and social responsibility at the center of our ambitions.

Sanofi is listed on EURONEXT: SAN and NASDAQ: SNY.

Regeneron Forward-Looking Statements and Use of Digital Media

This press release includes forward-looking statements that involve risks and uncertainties relating to future events and the future performance of Regeneron Pharmaceuticals, Inc. ("Regeneron" or the "Company"), and actual events or results may differ materially from these forward-looking statements. Words such as "anticipate," "expect," "intend," "plan," "believe," "seek," "estimate," variations of such words, and similar expressions are intended to identify such forward-looking statements, although not all forward-looking statements contain these identifying words. These statements concern, and these risks and uncertainties include, among others, the nature, timing, and possible success and therapeutic applications of products marketed or otherwise commercialized by Regeneron and/or its collaborators or licensees (collectively, "Regeneron's Products") and product candidates being developed by Regeneron and/or its collaborators or licensees (collectively, "Regeneron's Product Candidates") and research and clinical programs now underway or planned, including without limitation Dupixent; the likelihood, timing, and scope of possible regulatory approval and commercial launch of Regeneron's Product Candidates and new indications for Regeneron's Products, such as Dupixent for the treatment of chronic obstructive pulmonary disease with evidence of type 2 inflammation as discussed in this press release as well as for the treatment of pediatric eosinophilic esophagitis, chronic spontaneous urticaria, chronic pruritus of unknown origin, bullous pemphigoid, and other potential indications; uncertainty of the utilization, market acceptance, and commercial success of Regeneron's Products and Regeneron's Product Candidates and the impact of studies (whether conducted by Regeneron or others and whether mandated or voluntary), including the studies discussed or referenced in this press release, on any of the foregoing or any potential regulatory approval of Regeneron's Products (such as Dupixent) and Regeneron's Product Candidates; the ability of Regeneron's collaborators, licensees, suppliers, or other third parties (as applicable) to perform manufacturing, filling, finishing, packaging, labeling, distribution, and other steps related to Regeneron's Products and Regeneron's Product Candidates; the ability of Regeneron to manage supply chains for multiple products and product candidates; safety issues resulting from the administration of Regeneron's Products (such as Dupixent) and Regeneron's Product Candidates in patients, including serious complications or side effects in connection with the use of Regeneron's Products and Regeneron's Product Candidates in clinical trials; determinations by regulatory and administrative governmental authorities which may delay or restrict Regeneron's ability to continue to develop or commercialize Regeneron's Products and Regeneron's Product Candidates, including without limitation Dupixent; ongoing regulatory obligations and oversight impacting Regeneron's Products, research and clinical programs, and business, including those relating to patient privacy; the availability and extent of reimbursement of Regeneron's Products from third-party payers, including private payer healthcare and insurance programs, health maintenance organizations, pharmacy benefit management companies, and government programs such as Medicare and Medicaid; coverage and reimbursement determinations by such payers and new policies and procedures adopted by such payers; competing drugs and product candidates that may be superior to, or more cost effective than, Regeneron's Products and Regeneron's Product Candidates; the extent to which the results from the research and development programs conducted by Regeneron and/or its collaborators or licensees may be replicated in other studies and/or lead to advancement of product candidates to clinical trials, therapeutic applications, or regulatory approval; unanticipated expenses; the costs of developing, producing, and selling products; the ability of

Regeneron to meet any of its financial projections or guidance and changes to the assumptions underlying those projections or guidance; the potential for any license, collaboration, or supply agreement, including Regeneron's agreements with Sanofi and Bayer (or their respective affiliated companies, as applicable) to be cancelled or terminated; the impact of public health outbreaks, epidemics, or pandemics (such as the COVID-19 pandemic) on Regeneron's business; and risks associated with intellectual property of other parties and pending or future litigation relating thereto (including without limitation the patent litigation and other related proceedings relating to EYLEA® (afibercept) Injection, Praluent® (alirocumab), and REGEN-COV® (casirivimab and imdevimab)), other litigation and other proceedings and government investigations relating to the Company and/or its operations, the ultimate outcome of any such proceedings and investigations, and the impact any of the foregoing may have on Regeneron's business, prospects, operating results, and financial condition. A more complete description of these and other material risks can be found in Regeneron's filings with the U.S. Securities and Exchange Commission, including its Form 10-K for the year ended December 31, 2022 and its Form 10-Q for the quarterly period ended March 31, 2023. Any forward-looking statements are made based on management's current beliefs and judgment, and the reader is cautioned not to rely on any forward-looking statements made by Regeneron. Regeneron does not undertake any obligation to update (publicly or otherwise) any forward-looking statement, including without limitation any financial projection or guidance, whether as a result of new information, future events, or otherwise.

Regeneron uses its media and investor relations website and social media outlets to publish important information about the Company, including information that may be deemed material to investors. Financial and other information about Regeneron is routinely posted and is accessible on Regeneron's media and investor relations website (<http://newsroom.regeneron.com>) and its Twitter feed (<http://twitter.com/regeneron>).

Sanofi Disclaimers or Forward-Looking Statements

This press release contains forward-looking statements as defined in the Private Securities Litigation Reform Act of 1995, as amended. Forward-looking statements are statements that are not historical facts. These statements include projections and estimates regarding the marketing and other potential of the product, or regarding potential future revenues from the product. Forward-looking statements are generally identified by the words "expects", "anticipates", "believes", "intends", "estimates", "plans" and similar expressions. Although Sanofi's management believes that the expectations reflected in such forward-looking statements are reasonable, investors are cautioned that forward-looking information and statements are subject to various risks and uncertainties, many of which are difficult to predict and generally beyond the control of Sanofi, that could cause actual results and developments to differ materially from those expressed in, or implied or projected by, the forward-looking information and statements. These risks and uncertainties include among other things, unexpected regulatory actions or delays, or government regulation generally, that could affect the availability or commercial potential of the product, the fact that product may not be commercially successful, the uncertainties inherent in research and development, including future clinical data and analysis of existing clinical data relating to the product, including post marketing, unexpected safety, quality or manufacturing issues, competition in general, risks associated with intellectual property and any related future litigation and the ultimate outcome of such litigation, and volatile economic and market conditions, and the impact that pandemics or other global crises may have on us, our customers, suppliers, vendors, and other business partners, and the financial condition of any one of them, as well as on our employees and on the global economy as a whole. The risks and uncertainties also include the uncertainties discussed or identified in the public filings with the SEC and the AMF made by Sanofi, including those listed under "Risk Factors" and "Cautionary Statement Regarding Forward-Looking Statements" in Sanofi's annual report on Form 20-F for the year ended December 31, 2022. Other than as required by applicable law, Sanofi does not undertake any obligation to update or revise any forward-looking information or statements.

Regeneron Contacts:

Media Relations

Sharon Chen

Tel: +1 914-847-1546

Sharon.Chen@regeneron.com

Investor Relations

Vesna Tosic

Tel: +1 914-847-5443

Vesna.Tosic@regeneron.com

Sanofi Contacts:

Media Relations

Sally Bain

Tel: +1 617-834-6026

Sally.Bain@sanofi.com

Investor Relations

Eva Schaefer-Jansen

Tel: +33 7 86 80 56 39

eva.schaefer-jansen@sanofi.com

Arnaud Delepine

Tel: +33 (0)6 73 69 36 93

arnaud.delepine@sanofi.com

Corentine Driancourt

Tel: +33 (0)6 40 56 92 21

corentine.driancourt@sanofi.com

Felix Lauscher

Tel: +1 908-612-7239

felix.lauscher@sanofi.com

Tarik Elgoutni

Tel: +1 617-710-3587

tarik.elgoutni@sanofi.com

Nathalie Pham

Tel: +33 (0)7 85 93 30 17

nathalie.pham@sanofi.com

REGENERON

Source: Regeneron Pharmaceuticals, Inc.