



## Dupixent® (dupilumab) Demonstrates Improved Esophageal Function in Eosinophilic Esophagitis (EoE) Phase 4 Trial

May 5, 2026 at 7:00 AM EDT

**Dupixent showed significant and clinically meaningful improvements in both esophageal distensibility as well as disease-related structural changes and inflammation in the esophagus in adult patients with EoE at week 24 compared to placebo, in results presented at DDW**

**These results reinforce the roles of IL-4 and IL-13, two of the key and central drivers of type 2 inflammation, in EoE**

**EoE is a chronic, progressive disease that causes scarring and narrowing of the esophagus, making it difficult to swallow food, with more pronounced damage increasing the risk for severe symptoms**

TARRYTOWN, N.Y. and PARIS, May 05, 2026 (GLOBE NEWSWIRE) -- Regeneron Pharmaceuticals, Inc. (NASDAQ: REGN) and Sanofi today presented results from the REMODEL Phase 4 trial demonstrating that Dupixent® (dupilumab) showed significant and clinically meaningful improvements in both esophageal distensibility (a measure of esophageal function) as well as disease-related structural changes to the esophagus compared to placebo in adults with eosinophilic esophagitis (EoE) at week 24. The results reinforce the previously established ability of Dupixent to address underlying inflammation and symptoms of this disease; Dupixent was the first and is still the only biologic and leading treatment indicated for EoE. The new data were shared today in a late-breaking oral presentation at the 2026 Digestive Disease Week (DDW) conference.

EoE is a chronic, progressive inflammatory disease associated with type 2 inflammation that damages the esophagus and prevents it from working properly. For people with EoE, swallowing even small amounts of food can lead to choking and become a painful and worrisome experience. They are often left to contend with the frustration and anxiety of a constantly evolving list of foods to avoid, a poor quality of life and a higher risk of depression. In cases where EoE causes the esophagus to narrow, and the esophagus can no longer distend to support normal swallowing, forced and potentially painful dilation (physical expansion) of the esophagus may be needed. In severe cases, a feeding tube was historically the only available option to ensure proper caloric intake and adequate nutrition. However, Dupixent has dramatically changed the standard of care for this disease, improving quality of life for many patients.

“For patients who have eosinophilic esophagitis, if the disease remains untreated, it can progress to narrowing of the esophagus, more severe symptoms, such as food impaction, and the need for esophageal dilation,” said Evan S. Dellon, M.D., M.P.H., Professor of Gastroenterology and Hepatology at the University of North Carolina School of Medicine and lead author of this study. “Dupixent showed a potential to modify the course of eosinophilic esophagitis – by improving esophageal size at just 6 months – the magnitude of this improvement corresponded to the benefit that could be seen from an esophageal dilation procedure. It also reduced hallmark endoscopic and histologic signs of disease, which further strengthens the evidence that type 2 inflammation plays an important role in the biology of this disease. The ongoing study will allow us to learn the even longer-term impact of Dupixent on the scarring and narrowing of the esophagus in EoE.”

In the REMODEL Phase 4 trial, 69 adults with EoE were treated with Dupixent 300 mg (n=46) every week or placebo (n=23). At week 24, the results for Dupixent compared to placebo were:

- **Improved esophageal distensibility (a measure of esophageal function):** 1.28 mm improvement in esophageal distensibility plateau from baseline compared to -0.01 mm (1.30 mm placebo-corrected improvement;  $p < 0.05$ ), the primary endpoint. This represents a 9% improvement from baseline compared to 1% (8% placebo-corrected improvement;  $p < 0.05$ ), the key secondary endpoint. The esophageal distensibility plateau measures the ability of the esophagus to expand and allow food to be transported through the esophagus effectively, with higher numbers representing better esophageal function and lower numbers representing worse function.
- **Reduced disease-related structural changes to the esophagus:** 4.89-point reduction from baseline in abnormal endoscopic findings, as assessed by EREFS (a scale ranging from 0-18), evaluating a combination of esophageal edema (swelling), rings (bands of scar tissue that narrow the esophagus), exudates (white plaques indicative of inflammation), furrows (vertical grooves indicative of damaged lining) and strictures (narrowing), compared to 0.07-point increase (4.96-point placebo-corrected reduction;  $p < 0.0001$ ).
- **Improved histological findings:** 0.89-point and 0.80-point reductions from baseline in disease severity and extent, respectively, as assessed by the EoE-HSS grade and stage scores (both scales ranging from 0-3), measuring changes in eight cellular and tissue features of biopsy specimens at the microscopic level, compared to 0.18-point and 0.14-point reductions, respectively (0.71-point and 0.65-point placebo-corrected reductions, respectively;  $p < 0.0001$  for both).
- **Increased histological remission:** 59% of patients achieved peak esophageal intraepithelial counts of  $\leq 6$  eosinophils per

high-power field (eos/hpf) compared to 4% ( $p < 0.0001$ ).

- Additionally, 78% of patients achieved peak esophageal intraepithelial counts of  $< 15$  eos/hpf, the diagnostic threshold for EoE, compared to 4% (nominal  $p < 0.0001$ ).

The safety results from the REMODEL trial were generally consistent with the known safety profile of Dupixent in EoE. The overall rates of adverse events (AEs) were 62% for Dupixent and 48% for placebo. AEs more commonly observed with Dupixent than placebo included injection site pain (9% vs. 4%) and headache (9% vs. 4%). There were no serious AEs in either treatment group.

### **About REMODEL**

The ongoing REMODEL Phase 4, randomized, double-blind, placebo-controlled trial is evaluating the impact of Dupixent on esophageal remodeling and function in adults with EoE. During the 24-week, double-blind treatment period, all patients received Dupixent 300 mg every week or placebo. After week 24, patients entered an open-label treatment period and could either continue Dupixent or switch from placebo to Dupixent through week 128.

The primary endpoint assessed change from baseline at week 24 in esophageal distensibility plateau as measured by endoluminal functional lumen imaging probe (EndoFLIP), an innovative approach to assess esophageal function based on scarring and narrowing. Further assessments of esophageal distensibility are planned at week 76 and week 128 during the open-label treatment period. Esophageal distensibility plateau is reported in millimeters and represents how the esophagus expands in response to a small balloon being inflated at multiple points inside the esophagus. The distensibility plateau is the maximum esophageal diameter achieved at the narrowest point of the esophagus when further increases in balloon pressure do not meaningfully increase diameter. Higher numbers represent better esophageal function, and lower numbers represent worse function.

Secondary endpoints assessed at week 24 included:

- Percent change from baseline in esophageal distensibility plateau as measured by EndoFLIP, the key secondary endpoint.
- Change from baseline in EoE Endoscopic Reference Score (EoE-EREFs) on a 0-18 scale.
- Change from baseline in EoE Histology Scoring System (EoE-HSS) grade and stage scores, which measure changes in eight cellular and tissue features on 0-3 scales, respectively.
- Proportion of patients achieving histological disease remission (peak esophageal intraepithelial eosinophil count of  $\leq 6$  eosinophils [eos]/high power field [hpf]).
- Proportion of patients achieving the diagnostic threshold for EoE (peak esophageal intraepithelial eosinophil count of  $< 15$  eos/hpf).

### **About Dupixent**

Dupixent, which was invented using Regeneron's proprietary *VelocImmune*<sup>®</sup> technology, is a fully human monoclonal antibody that inhibits the signaling of the interleukin-4 (IL-4) and interleukin-13 (IL-13) pathways and is not an immunosuppressant. The Dupixent development program has shown significant clinical benefit and a decrease in type 2 inflammation in Phase 3 trials, establishing that IL-4 and IL-13 are two of the key and central drivers of the type 2 inflammation that plays a major role in multiple related and often co-morbid diseases.

Dupixent has received regulatory approvals in more than 60 countries in one or more indications including certain patients with atopic dermatitis, asthma, chronic rhinosinusitis with nasal polyps (CRSwNP), EoE, prurigo nodularis, chronic spontaneous urticaria (CSU), chronic obstructive pulmonary disease (COPD), bullous pemphigoid (BP) and allergic fungal rhinosinusitis (AFRS) in different age populations. More than 1,400,000 patients are being treated with Dupixent globally.<sup>1</sup>

### **About Regeneron's *VelocImmune* Technology**

Regeneron's *VelocImmune* technology utilizes a proprietary genetically engineered mouse platform endowed with a genetically humanized immune system to produce optimized fully human antibodies. When Regeneron's co-Founder, Board co-Chair, President and Chief Scientific Officer George D. Yancopoulos was a graduate student with his mentor Frederick W. Alt in 1985, they were the first to [envision](#) making such a genetically humanized mouse, and Regeneron has spent decades inventing and developing *VelocImmune* and related *VelociSuite*<sup>®</sup> technologies. Dr. Yancopoulos and his team have used *VelocImmune* technology to create a substantial proportion of all original, FDA-approved fully human monoclonal antibodies. This includes Dupixent<sup>®</sup> (dupilumab), Libtayo<sup>®</sup> (cemiplimab-rwlc), Praluent<sup>®</sup> (alirocumab), Kevzara<sup>®</sup> (sarilumab), Evkeeza<sup>®</sup> (evinacumab-dgng), Inmazeb<sup>®</sup> (atoltivimab, maftivimab and odesivimab-ebng) and Veopoz<sup>®</sup> (pozelimab-bbfg). In addition, REGEN-COV<sup>®</sup> (casirivimab and imdevimab) had been authorized by the FDA during the COVID-19 pandemic until 2024.

### **Dupilumab Development Program**

Dupilumab is being jointly developed by Regeneron and Sanofi under a global collaboration agreement. To date, dupilumab has been studied across more than 60 clinical trials involving more than 12,000 patients with various chronic diseases driven in part by type 2 inflammation.

In addition to the currently approved indications, Regeneron and Sanofi are studying dupilumab in a broad range of diseases driven by type 2 inflammation or other allergic processes in Phase 3 trials, including chronic pruritus of unknown origin and lichen simplex chronicus. These potential uses of dupilumab are currently under clinical investigation, and the safety and efficacy in these conditions have not been fully evaluated by any regulatory authority.

## U.S. INDICATIONS

DUPIXENT is a prescription medicine used:

- to treat adults and children 6 months of age and older with moderate-to-severe eczema (atopic dermatitis or AD) that is not well controlled with prescription therapies used on the skin (topical), or who cannot use topical therapies. DUPIXENT can be used with or without topical corticosteroids. It is not known if DUPIXENT is safe and effective in children with AD under 6 months of age.
- with other asthma medicines for the maintenance treatment of moderate-to-severe eosinophilic or oral steroid dependent asthma in adults and children 6 years of age and older whose asthma is not controlled with their current asthma medicines. DUPIXENT helps prevent severe asthma attacks (exacerbations) and can improve your breathing. DUPIXENT may also help reduce the amount of oral corticosteroids you need while preventing severe asthma attacks and improving your breathing. It is not known if DUPIXENT is safe and effective in children with asthma under 6 years of age.
- with other medicines for the maintenance treatment of chronic rhinosinusitis with nasal polyps (CRSwNP) in adults and children 12 years of age and older whose disease is not controlled. It is not known if DUPIXENT is safe and effective in children with CRSwNP under 12 years of age.
- to treat adults and children 1 year of age and older with eosinophilic esophagitis (EoE), who weigh at least 33 pounds (15 kg). It is not known if DUPIXENT is safe and effective in children with EoE under 1 year of age, or who weigh less than 33 pounds (15 kg).
- to treat adults with prurigo nodularis (PN). It is not known if DUPIXENT is safe and effective in children with PN under 18 years of age.
- with other medicines for the maintenance treatment of adults with inadequately controlled chronic obstructive pulmonary disease (COPD) and a high number of blood eosinophils (a type of white blood cell that may contribute to your COPD). DUPIXENT is used to reduce the number of flare-ups (the worsening of your COPD symptoms for several days) and can improve your breathing. It is not known if DUPIXENT is safe and effective in children with COPD under 18 years of age.
- to treat adults and children 2 years of age and older with chronic spontaneous urticaria (CSU) who continue to have hives that are not controlled with H1 antihistamine treatment. It is not known if DUPIXENT is safe and effective in children with CSU under 2 years of age, or who weigh less than 11 pounds (5 kg).
- to treat adults with bullous pemphigoid (BP). It is not known if DUPIXENT is safe and effective in children with BP under 18 years of age.
- to treat adults and children 6 years of age and older with allergic fungal rhinosinusitis (AFRS), who have had surgery on their nose or sinuses in the past. It is not known if DUPIXENT is safe and effective in children with AFRS under 6 years of age.

**DUPIXENT is not used** to relieve sudden breathing problems and will not replace an inhaled rescue medicine **or** to treat any other forms of hives (urticaria).

## IMPORTANT SAFETY INFORMATION

**Do not use** if you are allergic to dupilumab or to any of the ingredients in DUPIXENT®.

**Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:**

- have eye problems.
- have a parasitic (helminth) infection.
- are scheduled to receive any vaccinations. You should not receive a “live vaccine” right before and during treatment with DUPIXENT.
- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.
  - A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby.
- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Especially tell your healthcare provider if you** are taking oral, topical, or inhaled corticosteroid medicines; have asthma and use an asthma medicine; or have AD, CRSwNP, EoE, PN, COPD, CSU, BP, or AFRS and also have asthma. **Do not** change or stop your other medicines, including corticosteroid medicine or other asthma medicine, without talking to your healthcare provider. This may cause other symptoms that were controlled by those medicines to come back.

**DUPIXENT can cause serious side effects, including:**

- **Allergic reactions. DUPIXENT can cause allergic reactions, including skin reactions, that can sometimes be severe.** Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, skin rash, including rash that looks like a bullseye, painful red or blue bumps under the skin, or red pus-filled spots on the skin, general ill feeling, itching, swollen lymph nodes, nausea or vomiting, joint pain, or cramps in your stomach area.

- **Eye problems.** Tell your healthcare provider right away if you have any new or worsening eye problems, including eye pain or changes in vision, such as blurred vision. Your healthcare provider may send you to an ophthalmologist for an exam if needed.
- **Inflammation of your blood vessels.** Rarely, this can happen in people with asthma who receive DUPIXENT. This may happen in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered. Tell your healthcare provider right away if you get: rash, chest pain, worsening shortness of breath, brown or dark colored urine, persistent fever, or a feeling of pins and needles or numbness of your arms or legs.
- **Psoriasis.** This can happen in people with atopic dermatitis and asthma who receive DUPIXENT. Tell your healthcare provider about any new skin symptoms. Your healthcare provider may send you to a dermatologist for an examination if needed.
- **Joint aches and pain.** Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

#### The most common side effects:

- **Eczema:** injection site reactions; eye problems, including eye and eyelid inflammation, redness, swelling, itching, eye infection, dry eye, and blurred vision; cold sores in your mouth or on your lips; and high count of a certain white blood cell (eosinophilia).
- **Asthma:** injection site reactions; high count of a certain white blood cell (eosinophilia); pain in the throat (oropharyngeal pain); and parasitic (helminth) infections.
- **Chronic Rhinosinusitis with Nasal Polyps:** injection site reactions; eye problems, including eye and eyelid inflammation, redness, swelling, itching, eye infection, and blurred vision; high count of a certain white blood cell (eosinophilia), stomach problems (gastritis); joint pain (arthralgia); trouble sleeping (insomnia); and toothache.
- **Eosinophilic Esophagitis:** injection site reactions; upper respiratory tract infections; cold sores in your mouth or on your lips; and joint pain (arthralgia).
- **Prurigo Nodularis:** eye problems, including eye and eyelid inflammation, redness, swelling, itching, and blurred vision; herpes virus infections; common cold symptoms (nasopharyngitis); dizziness; muscle pain; and diarrhea.
- **Chronic Obstructive Pulmonary Disease:** injection site reactions; common cold symptoms (nasopharyngitis); high count of a certain white blood cell (eosinophilia); viral infection; back pain; inflammation inside the nose (rhinitis); diarrhea; stomach problems (gastritis); joint pain (arthralgia); toothache; headache; and urinary tract infection.
- **Chronic Spontaneous Urticaria:** injection site reactions.
- **Bullous Pemphigoid:** joint pain (arthralgia); eye problems, including eye and eyelid inflammation, redness, swelling, itching, and blurred vision; and herpes virus infections.
- **Allergic Fungal Rhinosinusitis:** injection site reactions; eye problems, including eye and eyelid inflammation, redness, swelling, itching, eye infection, and blurred vision; high count of a certain white blood cell (eosinophilia); stomach problems (gastritis); joint pain (arthralgia); trouble sleeping (insomnia); and toothache.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 6 months to less than 12 years of age, DUPIXENT should be given by a caregiver.

Please see accompanying full [Prescribing Information](#) including Patient Information.

#### About Regeneron

Regeneron (NASDAQ: REGN) is a leading biotechnology company that invents, develops and commercializes life-transforming medicines for people with serious diseases. Founded and led by physician-scientists, our unique ability to repeatedly and consistently translate science into medicine has led to numerous approved treatments and product candidates in development, most of which were homegrown in our laboratories. Our medicines and pipeline are designed to help patients with eye diseases, allergic and inflammatory diseases, cancer, cardiovascular and metabolic diseases, neurological diseases, hematologic conditions, infectious diseases, and rare diseases.

Regeneron pushes the boundaries of scientific discovery and accelerates drug development using our proprietary technologies, such as *VelociSuite*, which produces optimized fully human antibodies and new classes of bispecific antibodies. We are shaping the next frontier of medicine with data-powered insights from the Regeneron Genetics Center® and pioneering genetic medicine platforms, enabling us to identify innovative targets and complementary approaches to potentially treat or cure diseases.

For more information, please visit [www.Regeneron.com](http://www.Regeneron.com) or follow Regeneron on [LinkedIn](#), [Instagram](#), [Facebook](#) or [X](#).

#### About Sanofi

Sanofi is an R&D driven, AI-powered biopharma company committed to improving people's lives and delivering compelling growth. We apply our deep understanding of the immune system to invent medicines and vaccines that treat and protect millions of people

around the world, with an innovative pipeline that could benefit millions more. Our team is guided by one purpose: we chase the miracles of science to improve people's lives; this inspires us to drive progress and deliver positive impact for our people and the communities we serve, by addressing the most urgent healthcare, environmental, and societal challenges of our time.

Sanofi is listed on Euronext: SAN and Nasdaq: Sny.

### **Regeneron Forward-Looking Statements and Use of Digital Media**

*This press release includes forward-looking statements that involve risks and uncertainties relating to future events and the future performance of Regeneron Pharmaceuticals, Inc. ("Regeneron" or the "Company"), and actual events or results may differ materially from these forward-looking statements. Words such as "anticipate," "expect," "intend," "plan," "believe," "seek," "estimate," variations of such words, and similar expressions are intended to identify such forward-looking statements, although not all forward-looking statements contain these identifying words. These statements concern, and these risks and uncertainties include, among others, the nature, timing, and possible success and therapeutic applications of products marketed or otherwise commercialized by Regeneron and/or its collaborators or licensees (collectively, "Regeneron's Products") and product candidates being developed by Regeneron and/or its collaborators or licensees (collectively, "Regeneron's Product Candidates") and research and clinical programs now underway or planned, including without limitation Dupixent® (dupilumab) for the treatment of eosinophilic esophagitis; uncertainty of the utilization, market acceptance, and commercial success of Regeneron's Products (such as Dupixent) and Regeneron's Product Candidates and the impact of studies (whether conducted by Regeneron or others and whether mandated or voluntary), including the studies discussed or referenced in this press release, on any of the foregoing; the likelihood, timing, and scope of possible regulatory approval and commercial launch of Regeneron's Product Candidates and new indications for Regeneron's Products, including Dupixent for the treatment of chronic pruritus of unknown origin, lichen simplex chronicus, and other potential indications; the ability of Regeneron's collaborators, licensees, suppliers, or other third parties (as applicable) to perform manufacturing, filling, finishing, packaging, labeling, distribution, and other steps related to Regeneron's Products and Regeneron's Product Candidates; the ability of Regeneron to manage supply chains for multiple products and product candidates and risks associated with tariffs and other trade restrictions; safety issues resulting from the administration of Regeneron's Products (such as Dupixent) and Regeneron's Product Candidates in patients, including serious complications or side effects in connection with the use of Regeneron's Products and Regeneron's Product Candidates in clinical trials; determinations by regulatory and administrative governmental authorities which may delay or restrict Regeneron's ability to continue to develop or commercialize Regeneron's Products and Regeneron's Product Candidates; ongoing regulatory obligations and oversight impacting Regeneron's Products, research and clinical programs, and business, including those relating to patient privacy; the availability and extent of reimbursement or copay assistance for Regeneron's Products from third-party payors and other third parties, including private payor healthcare and insurance programs, health maintenance organizations, pharmacy benefit management companies, and government programs such as Medicare and Medicaid; coverage and reimbursement determinations by such payors and other third parties and new policies and procedures adopted by such payors and other third parties; changes to drug pricing regulations and requirements and Regeneron's pricing strategy, including in connection with Regeneron's April 2026 agreements with the U.S. government; other changes in laws, regulations, and policies affecting the healthcare industry; competing products and product candidates (including biosimilar products) that may be superior to, or more cost effective than, Regeneron's Products and Regeneron's Product Candidates; the extent to which the results from the research and development programs conducted by Regeneron and/or its collaborators or licensees may be replicated in other studies and/or lead to advancement of product candidates to clinical trials, therapeutic applications, or regulatory approval; unanticipated expenses; the costs of developing, producing, and selling products; the ability of Regeneron to meet any of its financial projections or guidance and changes to the assumptions underlying those projections or guidance; the potential for any license, collaboration, or supply agreement, including Regeneron's agreements with Sanofi and Bayer (or their respective affiliated companies, as applicable), to be cancelled or terminated; the impact of public health outbreaks, epidemics, or pandemics on Regeneron's business; and risks associated with litigation and other proceedings and government investigations relating to the Company and/or its operations (including the pending civil proceedings initiated or joined by the U.S. Department of Justice and the U.S. Attorney's Office for the District of Massachusetts), risks associated with intellectual property of other parties and pending or future litigation relating thereto (including without limitation the patent litigation and other related proceedings relating to EYLEA® (afibercept) Injection), the ultimate outcome of any such proceedings and investigations, and the impact any of the foregoing may have on Regeneron's business, prospects, operating results, and financial condition. A more complete description of these and other material risks can be found in Regeneron's filings with the U.S. Securities and Exchange Commission, including its Form 10-K for the year ended December 31, 2025 and its Form 10-Q for the quarterly period ended March 31, 2026. Any forward-looking statements are made based on management's current beliefs and judgment, and the reader is cautioned not to rely on any forward-looking statements made by Regeneron. Regeneron does not undertake any obligation to update (publicly or otherwise) any forward-looking statement, including without limitation any financial projection or guidance, whether as a result of new information, future events, or otherwise.*

*Regeneron uses its media and investor relations website and social media outlets to publish important information about the Company, including information that may be deemed material to investors. Financial and other information about Regeneron is routinely posted and is accessible on Regeneron's media and investor relations website (<https://investor.regeneron.com>) and its LinkedIn page (<https://www.linkedin.com/company/regeneron-pharmaceuticals>).*

### **Sanofi Disclaimers or Forward-Looking Statements**

*This press release contains forward-looking statements within the meaning of applicable securities laws, including the Private Securities Litigation Reform Act of 1995, as amended.*

*Forward-looking statements are statements that are not historical facts. These statements include projections and estimates and*

their underlying assumptions regarding the marketing and other potential of the product; regarding potential future events and revenues from the product. Words such as “expect,” “anticipate,” “believe,” “intend,” “estimate,” “plan,” “can,” “contemplate,” “could,” “is designed to,” “may,” “might,” “potential,” “objective,” “attempt,” “target,” “project,” “strategy,” “strive,” “desire,” “predict,” “forecast,” “ambition,” “guideline,” “seek,” “should,” “will,” “goal,” or the negative of these and similar expressions are intended to identify forward-looking statements.

Although Sanofi’s management believes that the expectations reflected in such forward-looking statements are reasonable, investors are cautioned that forward-looking information and statements are subject to various risks and uncertainties, many of which are difficult to predict and generally beyond the control of Sanofi, that could cause actual results and developments to differ materially from those expressed in, or implied or projected by, the forward-looking information and statements. These risks, uncertainties and assumptions include among other things, unexpected regulatory actions or delays, or government regulation generally, that could affect the availability or commercial potential of the product, the fact that product may not be commercially successful; authorities’ decisions regarding whether and when to approve a product candidate; political pressure in the United States to mandate lower drug prices including “most favored nation” pricing for State Medicaid programs; the uncertainties inherent in research and development, including future clinical data and analysis of existing clinical data relating to the product, including post marketing, unexpected safety, quality or manufacturing issues; competition in general; risks associated with intellectual property and any related pending or future litigation and the ultimate outcome of such litigation, and volatile economic and market conditions, and the impact that global crises may have on us, our customers, suppliers, vendors, and other business partners, and the financial condition of any one of them, as well as on our employees and on the global economy as a whole. The risks and uncertainties also include the uncertainties discussed or identified in the public filings with the SEC and the French Markets Authority (AMF) made by Sanofi, including those listed under “Risk Factors” and “Cautionary Statement Regarding Forward-Looking Statements” in Sanofi’s annual report on Form 20-F for the year ended December 31, 2025 or contained in our periodic reports on Form 6-K. Other than as required by applicable law, Sanofi does not undertake any obligation to update or revise any forward-looking information or statements. In light of these risks, uncertainties and assumptions, you should not place undue reliance on any forward-looking statements contained herein.

All trademarks mentioned in this press release are the property of the Sanofi group except for VelociSuite and Regeneron Genetics Center.

**Regeneron Contacts:**

**Media Relations**

**Kailey Kilmartin**

Tel: +1 914-652-0679

[Kailey.Kilmartin@regeneron.com](mailto:Kailey.Kilmartin@regeneron.com)

**Investor Relations**

**Mark Hudson**

Tel: +1 914-847-3482

[Mark.Hudson@regeneron.com](mailto:Mark.Hudson@regeneron.com)

**Sanofi Contacts:**

**Media Relations**

**Sandrine Guendoul**

Tel: +33 6 25 09 14 25

[Sandrine.Guendoul@sanofi.com](mailto:Sandrine.Guendoul@sanofi.com)

**Investor Relations**

**Thomas Kudsk Larsen**

Tel: +44 7545 513 693

[Thomas.Larsen@sanofi.com](mailto:Thomas.Larsen@sanofi.com)

**Evan Berland**

Tel: +1 215-432-0234

[Evan.Berland@sanofi.com](mailto:Evan.Berland@sanofi.com)

**Alizé Kaisserian**

Tel: +33 6 47 04 12 11

[Alize.Kaisserian@sanofi.com](mailto:Alize.Kaisserian@sanofi.com)

**Léo Le Bourhis**

Tel: + 33 6 75 06 43 81

[leo.lebourhis@sanofi.com](mailto:leo.lebourhis@sanofi.com)

**Keita Browne**

Tel: +1 781-249-1766

[Keita.Browne@sanofi.com](mailto:Keita.Browne@sanofi.com)

**Victor Rouault**

Tel: +1 617-356-4751

[Victor.Rouault@sanofi.com](mailto:Victor.Rouault@sanofi.com)

**Nathalie Pham**

Tel: +33 7 85 93 30 17

[Nathalie.Pham@sanofi.com](mailto:Nathalie.Pham@sanofi.com)

**Timothy Gilbert**

Tel: +1 516-521-2929

[Timothy.Gilbert@sanofi.com](mailto:Timothy.Gilbert@sanofi.com)

**Nina Goworek**

Tel : +1 908-569-7086

[Nina.Goworek@sanofi.com](mailto:Nina.Goworek@sanofi.com)

**Léa Ubaldi**

Tel: +33 6 30 19 66 46

[Lea.Ubaldi@sanofi.com](mailto:Lea.Ubaldi@sanofi.com)

**Thibaud Châtelet**

Tel: +33 6 80 80 89 90

[Thibaud.Chatelet@sanofi.com](mailto:Thibaud.Chatelet@sanofi.com)

**Ekaterina Pesheva**

Tel: +1 410-926-6780

[Ekaterina.Pesheva@sanofi.com](mailto:Ekaterina.Pesheva@sanofi.com)

**Yun Li**

Tel: +33 6 84 00 90 72

[Yun.Li3@sanofi.com](mailto:Yun.Li3@sanofi.com)

---

<sup>1</sup> Data on File

**REGENERON**

Source: Regeneron Pharmaceuticals, Inc.